

# Infectious Diseases Trends in Ontario, 2023



Technical Notes  
December 2024

## Public Health Ontario

Public Health Ontario is an agency of the Government of Ontario dedicated to protecting and promoting the health of all Ontarians and reducing inequities in health. Public Health Ontario links public health practitioners, frontline health workers and researchers to the best scientific intelligence and knowledge from around the world.

Public Health Ontario provides expert scientific and technical support to government, local public health units and health care providers relating to the following:

- communicable and infectious diseases
- infection prevention and control
- environmental and occupational health
- emergency preparedness
- health promotion, chronic disease and injury prevention
- public health laboratory services

Public Health Ontario's work also includes surveillance, epidemiology, research, professional development and knowledge services. For more information, visit [publichealthontario.ca](https://publichealthontario.ca).

How to cite this document:

Ontario Agency for Health Protection and Promotion (Public Health Ontario). Infectious disease trends in Ontario, 2023: technical notes. Toronto, ON: King's Printer for Ontario; 2024.

© King's Printer for Ontario, 2024

## Disclaimer

This document was developed by Public Health Ontario (PHO). PHO provides scientific and technical advice to Ontario's government, public health organizations and health care providers. PHO's work is guided by the current best available evidence at the time of publication.

The application and use of this document is the responsibility of the user. PHO assumes no liability resulting from any such application or use.

This document may be reproduced without permission for non-commercial purposes only and provided that appropriate credit is given to PHO. No changes and/or modifications may be made to this document without express written permission from PHO.

## Contents

Data Sources .....	1
Reporting .....	1
Integrated Public Health Information System and Public Health Case and Contact Management Solution .....	1
Public Health Ontario Laboratory Data.....	2
National Comparator Data.....	2
Case Definitions .....	3
Case Classifications .....	3
Data Management .....	5
Reference Period.....	5
Case Ascertainment Criteria .....	7
Re-Infection and Co-Infections .....	7
Descriptive Measures.....	8
Case Counts.....	8
Rates .....	9
Geographic Distribution.....	10
Public Health Unit Allocation .....	10
Age Distribution .....	10
Sex.....	11
Monthly Incidence .....	11
Hospitalizations.....	11
Deaths .....	12
Organism Details.....	12
Immunization Status .....	13
Importation Status.....	13
Risk Factors .....	13
Resistance Types.....	14
Specimen Testing.....	14
Drug Resistance.....	14
Analysis .....	15

Data Limitations .....	15
Accuracy of Data .....	15
Small Counts .....	15
Under-Reporting .....	16
Duplicates .....	16
Missing Data (Data Not Reported by PHUs) .....	16
Suggested Citations.....	17
Citation for the Infectious Disease Trends in Ontario Tool.....	17
Citation for a Specific Disease.....	17
Generic Citation Format.....	17
Source Statement for a Graph .....	17
Generic Citation Format.....	17
Source Statement for a Map.....	18
Generic Citation Format.....	18
Source Statement for Downloaded Data.....	18
Generic Citation Format.....	18
References .....	19
Appendix A: Diseases of Public Health Significance and Reportable Classifications: Ontario, 2023 .....	21
Appendix B: Summary of Updates Made to the IDTO Tool .....	25

# Data Sources

---

## Reporting

In Ontario, over 70 diseases are specified as diseases of public health significance under [Ontario Regulation 135/18: Designation of Diseases](#) pursuant to the [Health Protection and Promotion Act \(HPPA\), R.S.O 1990](#).<sup>1,2</sup>

Health care providers, laboratories and other individuals (including school principals and superintendents of institutions) with a duty to report diseases specified under Regulation 135/18 must make such reports to the Medical Officer of Health in the local public health unit (PHU) within which they operate, as outlined in [Regulation 569](#).<sup>3</sup> PHUs provide case management services to individuals suspected of having a disease of public health significance in their jurisdiction in accordance with the HPPA, the [Ontario Public Health Standards](#) and the [Infectious Diseases Protocol](#).<sup>4,5</sup> Required case data are subsequently reported to the province through the integrated Public Health Information System (iPHIS). Up to June 1, 2024 COVID-19 was reported through the Public Health Case and Contact Management Solution (CCM).

## Integrated Public Health Information System and Public Health Case and Contact Management Solution

The main source for disease data in the [Infectious Disease Trends in Ontario](#) interactive tool, with the exception of COVID-19, is the integrated Public Health Information System (iPHIS). iPHIS is the electronic reporting system used by all local PHUs to report cases of diseases of public health significance in Ontario.<sup>5</sup>

The data presented in the Infectious Disease Trends in Ontario tool are based mainly on cases with individual records in iPHIS. Aggregate counts and case-level data are used for both *Clostridium* (now more commonly known as *Clostridioides difficile*) infection (CDI) outbreaks in hospitals and varicella (chickenpox).

The iPHIS data used in this tool were extracted on July 3, 2024, with the exception of Carbapenemase-producing Enterobacteriaceae (CPE) data which were extracted on July 10, 2024.

The Public Health Case and Contact Management Solution (CCM) is the electronic reporting system that was used by all local PHUs to report cases of COVID-19 in Ontario (Note: As of June 2, 2024 entry and reporting of specific COVID-19 data shifted back to iPHIS with a focus on entering COVID-19 cases with fatal outcomes and outbreaks). IDTO 2023 includes available COVID-19 data for the 2020 (i.e., January 15 – December 31, 2020), 2021, 2022 and 2023 calendar years. CCM COVID-19 data in this tool were extracted on June 27, 2024 at 1 p.m.

It is important to note that iPHIS is dynamic disease reporting system that allows for ongoing updates to data previously entered. As a result, data extracted from iPHIS represent a snapshot at the time of extraction and may differ from previous or subsequent reports.

## Public Health Ontario Laboratory Data

Laboratory detection and confirmation of diseases of public health significance often occur at Public Health Ontario laboratory. For some diseases, identification can take place at hospitals or private laboratories. In other instances, reference services and specialized testing of clinical specimens take place at other reference laboratories across Canada, including the National Microbiology Laboratory (NML) and these results are reported back to PHO. The main source for laboratory test data in the 2023 Infectious Disease Trends in Ontario interactive tool is PHO's Laboratory Information Management System (LIMS). The testing data for calculating percent positivity for chlamydia, gonorrhea, influenza, legionellosis and resistance data for tuberculosis (TB) were extracted from LIMS on July 24, 2024.

Population estimates up to 2023 were provided by Statistics Canada on June 28, 2024 and were used as the denominator to calculate overall, age-, sex-, and PHU-specific crude incidence rates, where applicable.

Live birth counts for Ontario were originally sourced through Service Ontario and obtained from IntelliHEALTH Ontario. IntelliHEALTH Ontario is a repository of health-related data that describes the population and delivery of health care services in Ontario. Live births were used as the denominator for calculating incidence rates for two neonatal and congenital diseases included in this tool (i.e., neonatal group B streptococcal disease, and ophthalmia neonatorum) because neonatal population counts (infants up to 28 days old) could not be determined from available Vital Statistics data. The live birth data available to PHO are current up to 2022; as such, incidence rates for neonatal diseases for 2023 are based on the 2022 live birth data. Data on live births were extracted from IntelliHEALTH Ontario on June 6, 2024.

## National Comparator Data

Comparator incidence rates for Canada are provided in the tool whenever available. These data were obtained directly from the [Notifiable Disease Charts](#) of the Public Health Agency of Canada (PHAC) website on July 29, 2024.<sup>6</sup>

For most diseases in this tool, comparator incidence rates for Canada are presented for the years 2014 to 2022 in trend over time graphs. Depending on the disease and when it became nationally notifiable, national incidence rates may not be available for all or a part of this period. As a result, comparisons between trends in provincial and national incidence rates are made only for years in which national incidence rates are provided. National rates are also not included for diseases where the data provided do not distinguish between different forms of the disease (e.g., infectious and non-infectious syphilis and acute and chronic hepatitis B) or between aggregate and individual data on cases (i.e., varicella, CDI). Differences between national and Ontario case definitions also exist and these differences should be considered when comparing trends in provincial and national incidence rates. A list of diseases that are nationally notifiable can be found on [PHAC's website](#).<sup>7</sup>

Incidence rates for Canada presented in this tool may differ from reports published by PHAC. Where such discrepancies exist, incidence rates published in more recent national reports supersede those in this tool.

# Case Definitions

---

[Appendix A](#) in this document lists diseases of public health significance and associated case classifications that were reportable in Ontario in 2023. Cases are classified in iPHIS and CCM according to the provincial surveillance case definitions in use at the time the case was identified. PHUs are responsible for ensuring that cases reported to the province meet the relevant case definition. The most recent provincial case definitions are available in [Appendix 1](#) of the Infectious Diseases Protocol.<sup>8</sup>

It is important to consider changes in provincial case definitions and associated case classifications (see [Case Classifications](#) section below) over time when interpreting disease trends presented in this tool. The changes have occurred over the years to reflect the changing epidemiology of infectious diseases and the use of more current laboratory diagnostic practices and technology. Changes to the reportable diseases case definitions from 1991 to 2016 are documented in the [Factors affecting Reportable Diseases in Ontario \(1991-2016\)](#) report and its accompanying [appendix](#).<sup>9,10</sup>

# Case Classifications

---

Unless otherwise stated, case counts presented in this tool include only the confirmed case classification. Reporting of probable cases is only required for some diseases, as specified in the provincial case definitions outlined in [Appendix 1](#) of the Infectious Diseases Protocol.<sup>8</sup>

Probable cases are included in the total counts presented in the Infectious Disease Trends in Ontario tool for select diseases: amebiasis, anaplasmosis, babesiosis, Lyme disease, mumps, pertussis, Powassan virus, invasive meningococcal disease (IMD), invasive *Haemophilus influenzae* (Hi) disease all types, and West Nile virus (WNV) illness. Reporting of probable cases for these diseases, with the exception of WNV illness was instituted following case definition changes in 2009 because some cases that previously met the confirmed case definition were subsequently required to be reported as probable. As a result, both confirmed and probable cases are included in total counts for these diseases within this tool to ensure valid comparisons over time (see Table 1 for specific dates).

For the vast majority of other diseases that were similarly impacted by the 2009 case definition changes, the impact on overall counts was negligible. As such, probable cases for these diseases are not routinely included in provincial counts.



**Table 1: Diseases that include probable cases in total counts**

Disease	Case classifications
Amebiasis	Probable cases of amebiasis have been included in total counts since January 1, 2009 owing to the change in interpretation of laboratory test results that previously reported the causative agent as <i>Entamoeba histolytica</i> / <i>E. dispar</i> with no distinction between the two. Cases with test results that do not differentiate between the non-pathogenic <i>E. dispar</i> and the pathogenic <i>E. histolytica</i> are now counted as probable, whereas they were previously counted as confirmed. The impact of this change was substantial. As a result, probable cases have been included in total counts since 2009 to ensure valid comparisons over time for amebiasis.
Anaplasmosis	Both confirmed and probable cases have been included in provincial counts since the disease became reportable in July 1, 2023.
Babesiosis	Both confirmed and probable cases have been included in provincial counts since the disease became reportable in July 1, 2023.
<i>Haemophilus influenzae</i> (Hi)	<p>Probable cases have been included in total counts since April 28, 2009 to facilitate trending over time.</p> <p>As of May 1, 2018, all types of invasive Hi disease were designated as a disease of public health significance in Ontario. Previously, only type b was reportable. Thus, serotype information should be considered when comparing trends over time.</p>
Invasive meningococcal disease (IMD)	Probable cases have been included in total counts since April 28, 2009 to facilitate interpretation of trends over time.
Lyme disease	The impact of the change in case definition was substantial given that probable cases reported since 2009 constituted a substantial proportion of total case counts. As a result, probable Lyme diseases cases have been included in total counts since January 1, 2009 in order to ensure valid comparisons over time.
Mumps	Probable cases have been included in total counts since April 28, 2009 to facilitate interpretation of trends over time.
Pertussis	<p>Probable cases have been included in total counts since April 28, 2009 to facilitate interpretation of trends over time.</p> <p>Changes to laboratory testing may also impact temporal trends; testing by polymerase chain reaction (PCR), a more sensitive diagnostic tool, was first implemented in 1998, followed by real-time PCR in 2005. Effective 2009, the minimum threshold used to determine a positive PCR result was increased, leading to a reduction in the number of positive cases of pertussis identified via PCR.</p>
Powassan virus	Both confirmed and probable cases have been included in provincial counts since the disease became reportable in July 1, 2023.
West Nile virus (WNV) illness	Both confirmed and probable cases have been included in provincial counts since the disease became reportable in 2003.

For **measles** and **rubella**, probable cases are excluded from the historical temporal trend despite being reportable at the provincial level. These diseases, along with congenital rubella syndrome (CRS), have been declared eliminated from Canada and strict criteria are required to confirm or rule out cases. Since 2012, these criteria have been entrenched in provincially-enhanced surveillance activities to document the elimination of measles and rubella, which may impact the interpretation of trends in the incidence of these diseases. Despite elimination, Ontario continues to have cases due to importation from parts of the world where these diseases remain endemic.

For **hepatitis B**, confirmed acute cases are captured under the Classification Description of confirmed in iPHIS, while confirmed chronic hepatitis B cases are captured under the Classification Description of carrier. When a case progresses from acute to chronic infection, PHUs create a chronic carrier case, in addition to the existing acute confirmed case. Therefore, counts of acute and chronic hepatitis B cases are not mutually exclusive and have not been summed, as this would result in double-counting of some cases.

Both **AIDS** and **HIV** cases are reported under the Disease field in iPHIS as HIV/AIDS. HIV cases that have not progressed to AIDS have both an Encounter Type and a Diagnosis Status of carrier. HIV cases that have progressed to AIDS have an updated Encounter Type of case and an updated Diagnosis Status of confirmed. To determine accurate counts, cases of HIV/AIDS with either an Encounter Type of carrier and a Diagnosis Status of carrier or an Encounter Type of case and a Diagnosis Status of confirmed are counted as HIV cases using the Encounter Date (the date the HIV encounter was reported). HIV/AIDS encounters with an Encounter Type of case and a Diagnosis Status of confirmed are counted as AIDS cases based on the Diagnosis Status Date (the date the case was diagnosed with AIDS). Therefore, counts of AIDS and HIV cases are not mutually exclusive and have not been summed, as this would result in double-counting of some cases.

## Data Management

---

### Reference Period

iPHIS allows the entry of multiple dates relevant to the case. In Ontario, cases of most diseases of public health significance are classified by time using the episode date, which is a field that estimates the symptom onset date of disease for a case. The episode date is determined using the following hierarchy of date fields in iPHIS:

1. Symptom Onset Date
2. Specimen Collection Date
3. Lab Test Date (date laboratory testing was performed)
4. Reported Date (date the case was reported to the PHU)

During data extraction, the earliest available date in the hierarchy is selected as the episode date for each case. For example, if an Onset Date is available for a case, it is selected as the episode date instead of the Specimen Collection Date and so on. In some situations, the episode dates captured can be much later than the actual date of symptom onset, which can result in larger margins of error when deriving the symptom onset date from the Reported Date or the Specimen Collection date.

Cases are attributed to a particular year based on their episode date. There are some exceptions to reporting of incident cases; for example, HIV, chronic hepatitis B, hepatitis C, TB, late latent syphilis and neurosyphilis are often undiagnosed for extended periods and their detection by public health is generally not indicative of the actual date the infection was acquired. Therefore in some instances, cases included in this tool for a particular year are individuals who acquired their infections in earlier years and the data represent rates of new diagnoses rather than rates of new infection.

HIV/AIDS and TB are not classified by time based on the Episode Date. For HIV, incident case counts are based on the Encounter Date (i.e., Reported Date), defined as the date a case became known to public health. AIDS and TB incident case counts are based on the Diagnosis Status Date and Diagnosis Date, which is the date of a case's diagnosis for AIDS and TB, respectively.

Aggregate varicella counts are attributed to a year based on the reporting year as entered in the Outbreak Reported Year Name. If the Outbreak Reported Year Name was incomplete, the year indicated in the Outbreak Name (users are instructed to name outbreaks using the year of report) was used to attribute the cases to a particular year. If the Outbreak Name did not contain a year in the specified format, the year the outbreak was classified (the Outbreak Classification Year Name) was used.

CDI outbreaks are allocated to onset year based on the onset date of the index case in the outbreak. Where onset date of the index case was missing, the date the outbreak was created in iPHIS was used.

CPE case counts are based on the earliest specimen collection date. Cases with missing specimen collection date were excluded. CPE outbreaks are counted based on report date.

In CCM, episode date is based on the best estimate of the date of disease onset. This date is calculated based on either the date of symptom onset, specimen collection/test date for the COVID-19 test, or the date the COVID-19 case was reported to public health (not the specimen collection/test date). While the COVID-19 cases in IDTO are attributed to a particular year using this episode date hierarchy, PHO's Ontario Respiratory Virus Tool (ORVT) uses the date the COVID-19 case was reported to public health to place the data in time. In addition, the ORVT presents COVID-19 cases by surveillance weeks which correspond to the [FluWatch](#) Public Health Agency of Canada (PHAC) influenza surveillance weeks. Aggregated counts in the ORVT are presented by the surveillance period (approximately September 1 of one year to August 31 of the following year). Therefore, total annual cases in the IDTO will differ from those derived from the ORVT.

- Unless otherwise specified, the Infectious Disease Trends in Ontario tool covers the period from 2014 to 2023 for most diseases. Some exceptions include: Data for Blastomycosis, CPE infection or colonization and *Echinococcus multilocularis* infection are presented from 2018 onwards. These diseases were designated as diseases of public health significance in Ontario on May 1, 2018.

- Data for SARS COV-2 (COVID-19) are presented from 2020 onwards. This disease was first detected in Ontario on January 25, 2020.
- Data for mpox are presented from May 2022 onwards. This disease was designated as a disease of public health significance in Ontario on June 16, 2022 though cases were reported to the province before this date. Case counts, rates, hospitalizations and deaths for 2022 are based on eight months of data.
- Data for anaplasmosis, babesiosis and Powassan virus are presented from July 2023 onwards. These diseases were designated as a diseases of public health significance in Ontario on July 1, 2023. Case counts, rates, hospitalizations and deaths for 2023 are based on six months of data.

## Case Ascertainment Criteria

This tool includes all confirmed (and probable, as applicable) cases of diseases of public health significance reported through CCM with an episode date from 2020 to 2023 and through iPHIS with an episode date from 2014 to 2023 (or other appropriate date fields, as noted above in the [Reference Period](#) section), with the following exclusions:

1. Cases who were not residents of Ontario at the time of diagnosis.
2. Cases reported with a Disposition Status of entered in error, does not meet definition, duplicate – do not use, or any variation on these values.
3. Adverse events following immunization (AEFIs), which are summarized in an online tool available on PHU's [vaccine safety webpage](#).<sup>11</sup>
4. Cases reported as acute flaccid paralysis, encephalitis, meningitis, food poisoning or severe acute respiratory syndrome (SARS).
5. Institutional and public hospital outbreaks of gastroenteritis (where the Aetiologic Agent was not *Clostridioides difficile* in a hospital) and respiratory illness.
6. Cases with a missing outbreak number in iPHIS (i.e., sporadic cases should also have a sporadic outbreak number assigned in iPHIS where applicable).

[Appendix A](#) in this document provides a list of diseases of public health significance in Ontario in 2023 and notes the diseases that are excluded from this tool.

## Re-Infection and Co-Infections

For many of the diseases of public health significance, immunity is not conferred following infection or wanes over time, resulting in continued susceptibility and potential for re-infection. It is assumed that cases representing re-infection, as opposed to relapse, were assessed by PHUs before entry into iPHIS based on several factors, including the time period between the two episodes and the incubation period for the disease in question. As a result, data for most diseases in this tool are assumed to be new episodes of a disease or true re-infections. Thus, a single person with more than one episode of the same disease in

a single year may contribute more than one case of a particular disease to the total provincial count for that year. For example, this may occur for individuals with chlamydia, COVID-19, gonorrhoea or salmonellosis. For diseases caused by *Salmonella*, co-infections with two different serotypes (e.g., *Salmonella* Typhimurium and *Salmonella* Hadar) are reported as two separate episodes of salmonellosis. In addition, co-infections with more than one aetiologic agent at the same time (e.g., *Mycobacterium tuberculosis* complex and HIV) are reported as two different episodes, one for each disease caused by the co-infecting agents.

## Descriptive Measures

---

The descriptive measures used throughout the tool to characterize the epidemiology of diseases of public health significance in Ontario are listed below.

### Case Counts

This measure refers to the number of confirmed (and probable, as applicable) cases of a disease reported during a specified time frame and within a sub-group (if applicable) that meet the case ascertainment criteria outlined above. The diseases outlined in Table 2 are exceptions that require additional ascertainment based on the listed criteria.

**Table 2: Additional ascertainment criteria for AIDS, CDI outbreaks in hospitals, CPE, influenza, syphilis, tuberculosis and varicella (chickenpox)**

Disease	Additional ascertainment criteria
AIDS	AIDS case counts are based on the first AIDS diagnosis in a patient.
CDI outbreaks in hospitals	On September 1, 2008, Ontario amended regulations to make CDI outbreaks in public hospitals reportable to PHUs under the HPPA. <sup>1,12</sup> CDI outbreaks in long-term care homes that are reported as institutional outbreaks of gastroenteritis were excluded. For outbreak-level analyses, where discrepancies were observed between reported CDI aggregate case counts and line listed cases for the outbreak, counts of cases (and deaths) were determined based on the higher number. For case-level analyses, only individual confirmed case records associated with confirmed CDI outbreaks in hospitals were included for demographic and risk factor analyses. Cases with a non-reportable classification (e.g., probable cases) were excluded.
CPE	Case counts include infection, colonization and unspecified. Where multiple reports with the same carbapenemase are entered in iPHIS for a client, only the first report is included.

---

Disease	Additional ascertainment criteria
Influenza	Toronto Public Health (since mid-March 2020) and Ottawa Public Health (since mid-December 2023) share aggregate influenza case data with PHO each week for inclusion in the ORVT. These cases may not be entered in iPHIS which means that laboratory-confirmed influenza case counts since April 2020 that are presented in IDTO are likely incomplete. For more complete seasonal laboratory-confirmed influenza case totals, refer to the ORVT.
Syphilis	Includes only infectious cases (i.e., primary, secondary, early latent and infectious neurosyphilis).
TB	Includes only active cases in counts of confirmed cases (i.e., latent TB infections are not included).
Varicella (chickenpox)	<p>Varicella is reported provincially as both individual and aggregate cases. For individually-reported cases, which typically represent the more severe spectrum of disease (e.g., laboratory-confirmed cases, hospitalized cases, cases with complications), only confirmed cases are included.</p> <p>Aggregate cases represent the total number of cases reported in a PHU jurisdiction broken down by predefined age groups. They do not contain individual case details and may include cases that have been entered as individual cases, as well as those that do not meet the criteria for individual case reporting. For aggregate cases, only cases with Outbreak Classification field was entered as confirmed were included.<sup>13</sup></p>

## Rates

In general, incidence rate is defined as the number of new cases of disease (or newly reported/diagnosed cases for some diseases) occurring in a specified time period. Throughout the tool, the term “Rate” refers to an annual rate (e.g., the number of cases observed for every 100,000 Ontarians per year), unless otherwise specified. Crude incidence rates were calculated by dividing the total case count in a year by the total number of people in the population in that year. As specified in the [Case Classifications](#) section above, the total case count for some diseases may include confirmed and probable cases. The formulas for calculating overall and population-specific rates used throughout the tool are noted below (using the example of rates presented in a specified time period per 100,000 population).

$$\frac{\text{Number of cases in specified time period and population}}{\text{Total number of people in that population during the same time period}} \times 100,000$$

- **Overall rate:** Number of all new cases in a specified time period divided by the Ontario population for that time period, multiplied by 100,000.
- **Group-specific rate:** Number of new cases in a sub-group (e.g., age group, sex, PHU) in a specified time period divided by the population for that sub-group for that time period, multiplied by 100,000.

- **Neonatal rate:** Number of new congenital or neonatal cases of a disease (cases occurring in infants up to 28 days old) in a specified time period divided by the total number of live births for that time period, multiplied by 100,000. (Note: Live births with unknown geographic area are included in the Ontario live birth counts for that time period.)

---

**Note:** When data are downloaded using this interactive tool, zero values to the right of the decimal are lost (e.g., 5.0 appears as 5). Please ensure when using these data that all rate values are expressed to one decimal place (e.g., a value of 5 should be transformed to 5.0).

---

## Geographic Distribution

A PHU is a geographic area served by a Board of Health with the authority to deliver public health programs. Case counts and rates by PHU are available under the Maps tab for all years available for a disease.

## Public Health Unit Allocation

### CASES

In iPHIS, allocation of case counts by geography is based on Diagnosing Health Unit (DHU). The DHU is the PHU where the case resided when first detected. It does not necessarily reflect the location of exposure or diagnosis. iPHIS Bulletin #13 provides additional detail on scenarios in which a PHU is considered the DHU.<sup>14</sup> In CCM, the Permanent Health Unit field is equivalent to the DHU in iPHIS. For simplicity, DHU is used to denote both Diagnosing Health Unit and Permanent Health Unit throughout this document and in the interactive tool.

Incidence rates by PHU presented in the Maps tab are grouped into five categories: a category representing the lowest incidence (often zero) and four strata determined by dividing the range between the lowest and the highest PHU incidence rate into four equal intervals. These rate ranges are determined based on the entire time period included in the tool for the disease (i.e., 2014 to 2023 for most diseases) and the same categories are applied to all years of data. Rate range categories change when different sex and/or age group options are selected.

### CDI AND CPE OUTBREAKS

Allocation of CDI cases and outbreaks is based on the Primary Health Unit, which is the PHU where the hospital reporting the outbreak is located. While CPE case counts are allocated based on DHU, CPE outbreaks are allocated based on the Primary Health Unit.

## Age Distribution

Age groups for most diseases are based on standard five- and 10-year age groupings. For vaccine-preventable diseases, age groups are constructed with consideration of the epidemiology of the disease, and in some cases, the age of recommended vaccination. Cases with an unknown date of birth or a calculated age of less than zero or greater than 120, were classified as having an unknown age. Cases of unknown age are included when calculating total counts and rates, but are excluded in the calculation of age-specific rates.

## Sex

This tool uses the terminology sex to reference the reported values for the gender field in iPHIS. The exception is mpox, which uses the terminology gender. Three values for sex (gender in the case of mpox) are derived from the data entered in iPHIS: Male, Female, and Did Not Specify Male or Female. Information from all three fields are combined when presenting total counts or rates. For sex- or gender-specific rates, only male and female data are presented.

For COVID-19, male/female information was sourced from the sex field in CCM and is intended to represent sex assigned at birth. On October 14, 2021, changes were made in CCM to enable reporting on the sex field where this data field is supplemented by archived male/female information previously entered in the gender field.

## Monthly Incidence

For selected diseases, the number of cases that occurred during each month is available as an option for the Trends graph. For CDI, the number of outbreaks each month is compared to the monthly average outbreak count for the previous five years.

## Hospitalizations

This measure refers to the number of cases that were reported as hospitalized due to their disease, at the time of data extraction. In this tool, a case is considered hospitalized if at least one hospital admission date was recorded and the hospital admission date occurred 60 days or less before the episode date or 90 days or less after the episode date. The intention of applying an interval is to exclude hospital admission dates that are outside the reasonable range from the episode date to be attributed to the disease. The interval was selected to serve as cut points that could be applied in a uniform fashion across all diseases while including the majority of hospital records that fall within a reasonable range of the episode date.

Hospitalizations for COVID-19 include all cases admitted to hospital (or that had their hospital stay extended) because of COVID-19. This includes cases that have been discharged from hospital, cases that are currently in hospital and intensive care unit (ICU) cases but excludes emergency room visits. Hospital admissions were identified by a reported hospital admission date or reported “Yes” for hospitalization/ICU. If hospital admission date is missing, then ICU admission date is used (if applicable). When there is no ICU admission date to serve as a proxy, then reported date is used. Hospital admission date refers to the first admission date recorded on the case record. Hospital service transfers (e.g., alternate level of care (ALC)) are not reflected in the hospital admission date.

Emergency room visits are not included in the number of reported hospitalizations. It should be noted that under-reporting of hospitalizations may occur in iPHIS and CCM, particularly if hospitalization occurred after completion of follow-up by the PHU.

Hospitalization status is not reported for all diseases in this tool and is available only for the years 2015 to 2023 for diseases extracted from iPHIS, with the exception of mpox (2022 onward), anaplasmosis and babesiosis (available for 2023 only). COVID-19 hospitalizations are available for years 2020 onward. Severity indicators (hospitalizations and deaths) for influenza are not routinely collected and recorded in iPHIS and should therefore be interpreted with caution.



## Deaths

This measure refers to the number of cases that were reported as having died due to their disease, at the time of data extraction. For most diseases included in this tool, a case is counted as having died if at least one fatal outcome was recorded at the case level and the type/cause of death is a value other than “reportable disease was unrelated to cause of death”. Cases that have multiple types/causes of death, are counted as fatal if there is at least one value in this field other than “reportable disease was unrelated to cause of death.”

For TB, any case with a Date of Death entered in iPHIS is counted as fatal, except when the only cause of death entered in the iPHIS case record is “reportable disease was unrelated to cause of death”. The criteria for TB are different from those of most other diseases because the TB module is set up differently in iPHIS and the general criteria could not be applied. It should be noted that variations in follow-up may exist among PHUs in determining outcomes for all reportable diseases, as well as how the deaths are entered in the type/cause of death fields in iPHIS.

For CDI outbreak associated cases reported prior to 2015, any confirmed cases that were reported with an outcome of fatal or that had a date of death that occurred during the outbreak period, are counted as a fatal case. For CDI outbreak associated cases reported in 2015 and onwards, any confirmed cases reported with an outcome of fatal are counted as a fatal case. All reported deaths are classified as “all-cause” and may or may not be directly attributable to CDI.

A fatal COVID-19 case is defined as a death resulting from a clinically compatible illness in a confirmed COVID-19 case, unless there is a clear alternative cause of death that cannot be related to COVID-19 (e.g., trauma). There should be no period of complete recovery between the illness and the reported death. Deaths are determined by using the Outcome and Type of Death fields in CCM. COVID-19 deaths are counted where the Outcome value is “Fatal” and the Type of Death value is not “DOPHS was unrelated to cause of death” or “Under PHU Review”. COVID-19 deaths are placed in time using the Date of Death field in CCM. If the date of death is missing, the outcome date field is used as a proxy.

Under-reporting of deaths may occur in iPHIS and CCM, particularly if deaths occurred after completion of follow-up by the PHU. Cases occurring in 2023 for whom treatment is ongoing or if the disease is chronic may become fatal at a point in time after the extraction of data; these deaths would not be reflected in this tool. Case fatality data in 2023 for hepatitis B, hepatitis C, HIV/AIDS and TB are likely to be impacted by this issue.

Deaths are not reported for all diseases in this tool and are available only for the years 2015 to 2023 for diseases extracted from iPHIS, with the exception of blastomycosis (2018 onward), COVID-19 (2020 onward), mpox (2022 onward), anaplasmosis and babesiosis (2023 only).

## Organism Details

For select diseases, the tool provides the number and proportion of cases that represent distinct variations of a specific species, subtype, serotype, serogroup or genotype of a pathogen that causes a disease, for the years 2015 to 2023.

## Immunization Status

Immunization status is reported for Hi, measles, IMD, mumps, rubella and tetanus for the years 2015 to 2023 and is determined through an assessment of immunization records entered in iPHIS. In the absence of any immunization records, cases designated as unimmunized in the risk factor section of iPHIS are classified as unimmunized. When a case has no immunization records or information about whether or not they are unimmunized in the risk factor section in iPHIS, the case is classified as having an unknown immunization status. Immunizations are reported for cases who have received at least one dose of a vaccine related to the relevant disease prior to their illness onset. Not all immunizations recorded are considered valid; for example, vaccines administered prior to the minimum indicated age or vaccines that do not protect against the disease-causing serotype/serogroup would be excluded from the analysis.

## Importation Status

Importation status is reported for cases of measles and rubella for the years 2015 to 2023. A case of measles or rubella is considered imported if the person travelled outside Canada 7 to 21 and 14 to 21 days prior to symptom onset for measles and rubella, respectively.<sup>15</sup> Importation status is determined through a review of information entered in the risk factor field in iPHIS. In addition, data sent to the Canadian Measles/Rubella Surveillance System as part of the national enhanced surveillance of measles and rubella were consulted to validate the travel information in iPHIS.

## Risk Factors

---

For CPE case and case-level analyses of CDI outbreaks in hospitals, the number and proportion of cases that reported each risk factor are provided. Only cases reporting at least one risk factor were included in the risk factor analyses. Cases may report more than one risk factor. Risk factors are reported for CDI starting in 2015.

For CPE cases, as result of changes in iPHIS field configurations, risk factor variables that captured the same or similar concepts were combined as follows:

1. The risk factor Hospitalization in Canada in the last 12 months includes cases reporting “Yes” to the following risk factors:
  - Previous hospitalization at the reporting hospital in the last 12 months
  - Hospitalization in Canada in the last 12 months
  - Other inpatient hospitalization in Canada in the last 12 months
2. The risk factor Received health care outside of Canada in the last 12 months includes cases reporting “Yes” to the following risk factors:
  - Medical/Surgical procedure outside of Canada in the last 12 months
  - Hospitalization outside of Canada in the last 12 months
3. For consistency, the assumption was made that if cases had indicated they had received health care outside of Canada as a risk factor (see above) then cases also travelled outside of Canada in the last 12 months.

## Resistance Types

Reported carbapenemases were extracted from the Subtype Name field. Carbapenemase categories include New Delhi metallo- $\beta$ -lactamase (NDM), oxacillinase-48 carbapenemase (OXA-48), *Klebsiella pneumoniae* carbapenemase (KPC), Verona integron-encoded metallo- $\beta$ -lactamase (VIM) and 'Other' for all other carbapenemases, such as *Serratia marcescens* enzyme (SME) or Guiana extended-spectrum enzyme (GES). A CPE case may have more than one carbapenemase reported, which is reported as a single episode of CPE with multiple carbapenemases (e.g., KPC and OXA-48).

## Specimen Testing

Specimen testing data from PHO were used to calculate the percent positivity for chlamydia, gonorrhoea, legionellosis and influenza. Unless otherwise stated, these calculations were based on specimens received and tested at PHO from January 1, 2023 to December 31, 2023.

For chlamydia and gonorrhoea, the number of positive, negative and total tests performed at PHO is provided. Only nucleic acid amplification tests (NAAT) from cervical, urine, urethral, rectal and pharyngeal specimen sites are included. Rectal and pharyngeal specimens have been accepted for NAAT since April 2018. These data do not include testing performed at private laboratories throughout the province, which conduct the majority of testing for both chlamydia and gonorrhoea in Ontario. Percent positivity for chlamydia is calculated as the number of specimens positive for *Chlamydia trachomatis* divided by the total number of specimens tested for this pathogen. Percent positivity for gonorrhoea is calculated as the number of specimens positive for *Neisseria gonorrhoeae* divided by the total number of specimens tested for this pathogen.

For legionellosis, the percent positivity and number of total patients tested at PHO are provided. The month associated with each test was calculated based on the date PHO received the specimen. Data represents unique patients tested for *Legionella* by urine antigen, PCR or culture. Percent positivity is calculated as the number of patients positive for *Legionella* divided by the total number of patients tested for this pathogen.

For influenza, the percent positivity and the total number of specimens tested at PHO for influenza A (H3 and H1N1 pdm09) and influenza B are provided. These data do not include testing performed at private laboratories throughout the province, which conduct the majority of testing for influenza in Ontario. Data represents unique specimens tested for influenza by multiplex PCRs and culture. The week associated with each specimen was calculated as Sunday to Saturday (FluWatch weeks) and was based on the date the specimen was received by PHO. Percent positivity is calculated as the number of specimens positive for influenza A or B divided by the total number of specimens tested for influenza.

## Drug Resistance

Drug resistance data for tuberculosis from PHO were used to provide the number of multidrug-resistant TB cases and extensively drug-resistant TB cases. The numbers were based on specimens received and tested at PHO from January 1, 2023 to December 31, 2023.

# Analysis

---

The data analyses and presentation for the online tool was completed using SAS 9.4 and custom JavaScript code. Identified differences in rates and counts from one period to another, between Ontario and Canada and between sub-groups are population based and hence not subjected to analysis for statistical significance.

## Data Limitations

---

### Accuracy of Data

PHO coordinates an annual data cleaning exercise with PHUs to review cases/events of public health significance that occurred in their jurisdiction for the purpose of improving iPHIS data quality. The 2020 annual data cleaning process was interrupted due to the COVID-19 pandemic. The annual initiative resumed for 2021 data. However, there may have been variations in the ability of public health units to participate in the process. As a result, there may be limitations related to the completeness of 2020 through to 2023 data and trends should be interpreted with caution.

Any data extracted from iPHIS or CCM, including the data used in this tool represent a snapshot at the time of extraction and may differ from previous or subsequent reports. Discrepancies in disease counts and rates provided in this tool and other published data may exist due to:

- Enhanced data cleaning for select analyses, such as the linkage of iPHIS and laboratory data and subsequent reconciliation in iPHIS
- Late reporting
- Local and/or provincial-led data cleaning initiatives
- Differences in data extraction dates

Where such variability exists, data provided in the most recent release of this tool, other PHO surveillance tools and products (e.g., [Quarterly Infectious Diseases Surveillance Report](#)) or published research may be a more appropriate source depending on how the methodology, data caveats and/or extraction dates align with the intended use of the data.<sup>16</sup>

### Small Counts

For some diseases, the observed variability in population-specific incidence rates should be interpreted with caution owing to small counts, which may be exacerbated by small denominators (population). For this reason, users of this tool should be aware that these rates may be unstable.

## Under-Reporting

Passive surveillance systems, such as iPHIS and CCM, which rely primarily on mandatory health care provider and laboratory reports of illness can be characterized by under-reporting of the true burden of illness. For diseases captured in iPHIS, case counts only represent known cases reported to PHUs and recorded in the system. The resulting degree of under-reporting may vary from disease to disease due to a variety of factors. Interpret surveillance results for DOPHS in 2020 through 2023 with caution due to changes in the availability of health care, health seeking behaviours, public health follow-up, and case entry during the COVID-19 pandemic and subsequent recovery period.

Asymptomatic individuals who are colonized with CPE are only identified and reported if they are screened at a health care facility. As a result, colonized cases captured by provincial surveillance are more likely to have chronic underlying medical conditions that predispose their access to health care while asymptomatic cases in the community are likely underrepresented. Currently, health care facilities are in the process of implementing CPE screening programs and robustness may vary. Given the heterogeneity in CPE screening practices among health care facilities in Ontario, colonizations are likely underreported. In addition, there is a potential for misclassification of colonizations and infections due to variability in the interpretation of symptomatic presentation.

The COVID-19 data represent only cases reported to public health units and recorded in CCM. As a result, all counts will be subject to underreporting due to a variety of factors, such as disease awareness and medical care seeking behaviours, which may depend on severity of illness, clinical practice, changes in laboratory testing, and reporting behaviours.

## Duplicates

The potential for duplicates exists, as duplicate sets were not identified and excluded unless they were resolved prior to data extraction either at the local or provincial level. Inclusion of duplicates would result in over-reporting. (Note: An exception was made for CCM data, and duplicates were removed for COVID-19.)

## Missing Data (Data Not Reported by PHUs)

Data quality (completeness) for some fields is lower than others. Hospitalization and death are underreported in iPHIS and CCM, with the degree of underreporting influenced by the severity of illness and associated outcomes (e.g., less underreporting if illness or outcomes are more severe) and the timing of the event (i.e., there is likely less underreporting if hospitalization or death occurs shortly after symptom onset or before investigation of the case by the PHU is completed). Underreporting of risk factors, immunization status and specific laboratory data (e.g., serotype, genotype) can also occur. In general, the degree of underreporting is influenced by a combination of factors, including incomplete follow-up of cases (e.g., case is not reachable), incomplete or late entry of data in iPHIS/CCM, and the occurrence of outcomes after follow-up has been completed. A high proportion of missing or incomplete data may result in conclusions or interpretations that are not representative of the underlying epidemiology of the disease.

# Suggested Citations

---

## Citation for the Infectious Disease Trends in Ontario Tool

Ontario Agency for Health Protection and Promotion (Public Health Ontario). Infectious disease trends in Ontario [Internet]. Toronto, ON: King's Printer for Ontario; c2024 [cited YYYY Mon DD]. Available from: <https://www.publichealthontario.ca/en/data-and-analysis/infectious-disease/reportable-disease-trends-annually>

### Citation Tips

- It is important to include a cited date in order to transparently reflect the currency of the data.
- Use the URL specific to the disease rather than the main report. URLs are not included for graphs or maps because the URLs are unique to each disease, not the specific selections made to generate a particular graph or map.
- The details about the data sources are available in the [Data Sources](#) section of this document.

## Citation for a Specific Disease

### Generic Citation Format

Author. Interactive tool name: specific disease [Internet]. Toronto, ON: King's Printer for Ontario; cYear [cited date]. Available from: URL

### Example

Ontario Agency for Health Protection and Promotion (Public Health Ontario). Infectious disease trends in Ontario: chlamydia [Internet]. Toronto, ON: King's Printer for Ontario; c2024 [cited 2024 Dec 16]. Available from: <https://www.publichealthontario.ca/en/data-and-analysis/infectious-disease/reportable-disease-trends-annually#/11>

## Source Statement for a Graph

### Generic Citation Format

Source: Author. Interactive tool name: specific title as it appears on the graph [Internet]. Toronto, ON: King's Printer for Ontario; cYear [cited date].

### Example

Source: Ontario Agency for Health Protection and Promotion (Public Health Ontario). Infectious disease trends in Ontario: gonorrhoea cases by age, for all sexes, in Ontario [Internet]. Toronto, ON: King's Printer for Ontario; c2024 [cited 2024 Dec 16].

## Source Statement for a Map

### Generic Citation Format

Source: Author. Interactive tool name: specific title as it appears on the map [Internet]. Toronto, ON: King's Printer for Ontario; cYear [cited date].

#### Example

Source: Ontario Agency for Health Protection and Promotion (Public Health Ontario). Infectious disease trends in Ontario: tuberculosis rates for all ages, by males, 2013 [Internet]. Toronto, ON: King's Printer for Ontario; c2024 [cited 2024 Dec 16].

## Source Statement for Downloaded Data

### Generic Citation Format

Source: Data source as extracted and/or received by author. Interactive tool name: specific title as it appears on the source graph or map or other tab [Internet]. Toronto, ON: King's Printer for Ontario; cYear [cited date].

#### Example

Source: Data sources as extracted and/or received by Ontario Agency for Health Protection and Promotion (Public Health Ontario). Infectious disease trends in Ontario: West Nile virus illness rates and cases for all ages, for all sexes, Ontario [Internet]. Toronto, ON: King's Printer for Ontario; c2024 [cited 2024 Dec 16].

# References

---

1. *Designation of Diseases*, O Reg 135/18. Available from: <https://www.ontario.ca/laws/regulation/180135>
2. *Health Protection and Promotion Act*, RSO 1990, c H.7. Available from: <http://www.ontario.ca/laws/statute/90h07>
3. *RRO 1990, Reg 569*. Available from: <http://www.ontario.ca/laws/regulation/900569>
4. Ontario. Ministry of Health; Ministry of Long-Term Care. Ontario public health standards: requirements for programs, services, and accountability. Effective: June 2021 [Internet]. Toronto, ON: Queen's Printer for Ontario; 2021 [modified 2021 Jun; cited 2023 Dec 12]. Available from: <https://files.ontario.ca/moh-ontario-public-health-standards-en-2021.pdf>
5. Ontario. Ministry of Health Long-Term Care, Population and Public Health Division. Infectious diseases protocol, 2023. Effective: July 2023 [Internet]. Toronto, ON: King's Printer for Ontario; 2023 [cited 2024 Oct 24]. Available from: <https://files.ontario.ca/moh-infectious-disease-protocol-en-2023.pdf>
6. Public Health Agency of Canada. Notifiable disease charts [Internet]. Ottawa, ON: Government of Canada; 2024 [modified 2024 Aug 28; cited 2024 Oct 28]. Available from: <https://diseases.canada.ca/notifiable/charts-list>
7. Public Health Agency of Canada. Case definitions: nationally notifiable diseases [Internet]. Ottawa, ON: Government of Canada; 2024 [modified 2024 Aug 28; cited 2024 Oct 28]. Available from: <https://diseases.canada.ca/notifiable/diseases-list>
8. Ontario. Ministry of Health. Ontario public health standards: requirements for programs, services, and accountability. Infectious disease protocol. Appendix 1, provincial case definitions and disease specific information [Internet]. Toronto, ON: King's Printer for Ontario; 2023 [updated 2023 Dec 30; cited 2023 Dec 30]. Available from: <https://www.ontario.ca/page/ontario-public-health-standards-requirements-programs-services-and-accountability#section-2>
9. Ontario Agency for Health Protection and Promotion (Public Health Ontario). Factors affecting reportable diseases in Ontario: case definition changes and associated trends 1991-2016 [Internet]. Toronto, ON: Queen's Printer for Ontario; 2018 [cited 2020 Nov 10]. Available from: <https://www.publichealthontario.ca/-/media/documents/F/2018/factors-reportable-diseases-ontario-1991-2016.pdf?la=en&hash=A10D37CEE72926746247664DEA6E8E503AFAE0B2>
10. Ontario Agency for Health Protection and Promotion (Public Health Ontario). Appendix: factors affecting reportable diseases in Ontario: case definition changes and associated trends in Ontario: 1991-2016 [Internet]. Toronto, ON: Queen's Printer for Ontario; 2018 [cited 2020 Nov 16]. Available from: <https://www.publichealthontario.ca/-/media/documents/A/2018/appendix-factors-reportable-diseases-ontario-1991-2016.pdf?la=en&hash=EB2D15578B8C35E7DDFD64D8B192C27AD2025443>
11. Ontario Agency for Health Protection and Promotion (Public Health Ontario). Vaccine safety [Internet]. Toronto, ON: Queen's Printer for Ontario; 2024 [updated 2024 Mar 01; cited 2024 Oct 15]. Available from: <https://www.publichealthontario.ca/en/health-topics/immunization/vaccine-safety>



12. *Specification of Communicable Diseases*, O Reg 558/91. Available from: <https://www.ontario.ca/laws/regulation/910558>
13. Ontario. Ministry of Health and Long-Term Care. iPHIS final outbreak summary: user guide, version 5. Toronto, ON: Queen's Printer for Ontario; 2008.
14. Ontario Agency for Health Protection and Promotion (Public Health Ontario). iPHIS bulletin #13: transferring client responsibility[Internet]. Toronto, ON: Queen's Printer for Ontario; 2006 [revised 2020 Mar; cited 2021 Dec 08]. Available from: <https://www.publichealthontario.ca/-/media/documents/i/2020/iphis-bulletin-client-responsibility.pdf>
15. Lim GH, Deeks SL, Fediurek J, Gubbay J, Crowcroft NS. Documenting the elimination of measles, rubella, and congenital rubella syndrome in Ontario: 2009-12. *Can Commun Dis Rep*. 2014;40(8):143-51. Available from: <http://www.phac-aspc.gc.ca/publicat/ccdr-rmtc/14vol40/dr-rm40-08/dr-rm40-08-surv-eng.php>
16. Ontario Agency for Health Protection and Promotion (Public Health Ontario). Infectious disease surveillance reports [Internet]. Toronto, ON: King's Printer for Ontario; 2024 [updated 2024 Oct 16; cited 2024 Oct 28]. Available from: <https://www.publichealthontario.ca/en/data-and-analysis/infectious-disease/infectious-diseases-monthly>

# Appendix A: Diseases of Public Health Significance and Reportable Classifications: Ontario, 2023

Table A1 summarizes the diseases of public health significance, as specified under Ontario [Regulation 135/18<sup>1</sup>](#) and amendments under the [Health Protection and Promotion Act<sup>2</sup>](#) and the associated reportable case classifications, as outlined in [Appendix 1](#) of the Infectious Diseases Protocol.

**Table A1: Diseases of public health significance and reportable case classifications in Ontario, 2023**

Disease of public health significance	Reportable case classifications
Acquired immunodeficiency syndrome (HIV/AIDS)	Confirmed
Acute Flaccid Paralysis (AFP)*,+	Confirmed
Adverse events following immunization (AEFIs)*	Confirmed
Amebiasis <sup>‡</sup>	Confirmed, probable
Anaplasmosis <sup>‡,§</sup>	Confirmed, probable
Anthrax	Confirmed, probable, suspect
Babesiosis <sup>‡,§</sup>	Confirmed, probable
Blastomycosis <sup>¶</sup>	Confirmed, probable
Botulism	Confirmed, probable, suspect
Brucellosis	Confirmed, probable
<i>Campylobacter</i> enteritis	Confirmed, probable
Carbapenemase-producing <i>Enterobacteriaceae</i> (CPE) infection or colonization <sup>¶</sup>	Confirmed
Chancroid	Confirmed, probable
Chickenpox (Varicella)	Confirmed
<i>Chlamydia trachomatis</i> infections	Confirmed, probable
Cholera	Confirmed, probable

Disease of public health significance	Reportable case classifications
<i>Clostridium difficile</i> infection (CDI) Outbreaks in Public Hospitals <sup>#</sup>	Confirmed
Creutzfeldt-Jakob disease, all types	Confirmed, probable, suspect
Cryptosporidiosis	Confirmed, probable
Cyclosporiasis	Confirmed, probable
Diphtheria	Confirmed, probable
<i>Echinococcus multilocularis</i> infection <sup>¶</sup>	Confirmed, probable
Encephalitis* <ul style="list-style-type: none"> <li>• Primary, viral</li> <li>• Post-infectious</li> <li>• Vaccine-related</li> <li>• Subacute sclerosing panencephalitis</li> <li>• Unspecified</li> </ul>	Confirmed, probable
Food poisoning, all causes*	Confirmed, probable, suspect
Gastroenteritis, institutional outbreaks*	Not applicable
Giardiasis, except asymptomatic cases	Confirmed, probable
Gonorrhoea	Confirmed, probable
Group A Streptococcal disease, invasive	Confirmed
Group B Streptococcal disease, neonatal	Confirmed, probable
<i>Haemophilus influenzae</i> disease, all types, invasive <sup>‡,¶</sup>	Confirmed, probable
Hantavirus pulmonary syndrome	Confirmed
Hemorrhagic fevers, including: <ul style="list-style-type: none"> <li>• Ebola virus disease</li> <li>• Marburg virus disease</li> <li>• Lassa Fever</li> <li>• Other viral causes including bunyaviruses, arenaviruses and flaviviruses</li> </ul>	Confirmed, probable
Hepatitis A	Confirmed, probable
Hepatitis B**	Confirmed, Chronic, probable
Hepatitis C	Confirmed

Disease of public health significance	Reportable case classifications
Influenza	Confirmed
Legionellosis	Confirmed, probable
Leprosy	Confirmed, probable
Listeriosis	Confirmed, probable
Lyme disease <sup>‡</sup>	Confirmed, probable
Measles	Confirmed, probable
Meningitis, acute* <ul style="list-style-type: none"> <li>• Bacterial</li> <li>• Viral</li> <li>• Other</li> </ul>	Confirmed, probable
Meningococcal disease, invasive <sup>‡</sup>	Confirmed, probable
Mumps <sup>‡</sup>	Confirmed, probable
Diseases caused by a novel coronavirus*, including Coronavirus Disease 2019 (COVID-19), Severe Acute Respiratory Syndrome (SARS) and Middle East Respiratory Syndrome (MERS)	Confirmed, presumptive confirmed, probable, laboratory-based case of reinfection (COVID-19 only), time-based case of reinfection (COVID-19 only)
Ophthalmia neonatorum	Confirmed, probable
Paralytic Shellfish Poisoning (PSP) <sup>†</sup>	Confirmed, probable
Paratyphoid fever	Confirmed, probable
Pertussis (whooping cough) <sup>‡</sup>	Confirmed, probable
Plague	Confirmed, probable
Pneumococcal disease, invasive	Confirmed
Poliomyelitis, acute	Confirmed
Powassan virus <sup>‡,§</sup>	Confirmed, probable
Psittacosis/Ornithosis	Confirmed, probable
Q-fever	Confirmed, probable
Rabies	Confirmed, probable
Respiratory infection outbreaks in institutions and public hospitals*	Not applicable

Disease of public health significance	Reportable case classifications
Rubella	Confirmed, probable
Rubella, congenital syndrome	Confirmed, probable
Salmonellosis	Confirmed, probable
SARS COV-2 disease (COVID-19) <sup>††</sup>	Confirmed, probable
Shigellosis	Confirmed, probable
Smallpox and other Orthopoxviruses including mpox <sup>**</sup> (monkeypox)	Confirmed, probable, suspect, person under investigation
Syphilis, infectious	Confirmed
Tetanus	Confirmed
Trichinosis	Confirmed, probable
Tuberculosis	Confirmed, suspect
Tularemia	Confirmed, probable
Typhoid fever	Confirmed, probable
Verotoxin-producing <i>E. coli</i> infection indicator conditions, including Haemolytic Uremic Syndrome	Confirmed, probable
West Nile Virus illness <sup>‡</sup>	Confirmed, probable
Yersiniosis	Confirmed, probable

**Source:** Ontario. Ministry of Health. Ontario public health standards: requirements for programs, services, and accountability. Infectious disease protocol. Appendix 1, provincial case definitions and disease specific information [Internet]. Toronto, ON: Kings’s Printer for Ontario; 2024 [updated 2023 Dec 15; cited 2023 Dec 30]. Available from: <https://www.ontario.ca/page/ontario-public-health-standards-requirements-programs-services-and-accountability#section-2>

\*Disease not included in this tool.

<sup>†</sup>These diseases became reportable on December 4, 2013.

<sup>‡</sup>Both confirmed and probable cases of these diseases are included in this tool, whereas only confirmed cases are included for other diseases.

<sup>§</sup>This disease became reportable on July 1, 2023.

<sup>¶</sup>Disease became designated as a disease of public health significance on May 1, 2018. For invasive *Haemophilus influenzae* disease, all types of Hi were designated as a disease of public health significance, as of May 1, 2018, prior to which only type b was reportable.

<sup>#</sup>*Clostridium difficile* infection is now more commonly referred to as *Clostridioides difficile* infection.

<sup>\*\*</sup>Chronic case (carrier) classification added in 2012.

<sup>††</sup>This disease became reportable on January 22, 2020.

<sup>†††</sup>This disease became reportable on June 16, 2022.

# Appendix B: Summary of Updates Made to the IDTO Tool

---

Following are the updates made to the [Infectious Disease Trends in Ontario](#) interactive tool. For more information on updates, please contact [data@oahpp.ca](mailto:data@oahpp.ca).

Description	Date
Added three new diseases <ul style="list-style-type: none"> <li>Anaplasmosis</li> <li>Babesiosis</li> <li>Powassan virus</li> </ul>	December 2024
Removed Local Health Integration Network (LHIN) from interactive maps	December 2024
Added new disease <ul style="list-style-type: none"> <li>Mpox</li> </ul>	December 2023
Modified the blastomycosis page to present death data	December 2023
Modified the congenital rubella syndrome (CRS) page to present an 'Overview' tab. Removed trends, map, hospitalizations, deaths, importation status and immunization status tabs	December 2023
Removed malaria and yellow fever information	December 2023
Added new data for 2022 for existing diseases; re-extracted 2013-2021 data and updated the tool	December 2023
Updated Table 2 in the technical notes to include ascertainment criteria about AIDS and influenza cases	May 2023
Added new data for 2021 for existing diseases; re-extracted 2012-20 data and updated the tool	September 2022
Added new data for 2020 for existing diseases; re-extracted 2011-19 data and updated the tool	September 2021
Added a new disease: <ul style="list-style-type: none"> <li>COVID-19</li> </ul>	September 2021
Added new data for 2019 for existing diseases; re-extracted 2010-18 data and updated the tool	September 2020

Description	Date
Updated the number of public health units to 34 due to the merger of Huron and Perth health units	September 2020
Added new data for 2018 for existing diseases with the exception of malaria and yellow fever; re-extracted 2005–17 data and updated the tool	November 2019
Added new diseases: <ul style="list-style-type: none"> <li>• Blastomycosis</li> <li>• Carbapenemase-producing <i>Enterobacteriaceae</i> (CPE) infection or colonization</li> <li>• <i>Echinococcus multilocularis</i> infection (EMI)</li> </ul>	November 2019
Modified the invasive <i>Haemophilus influenzae</i> , type b page to present data for invasive <i>Haemophilus influenzae</i> , all types from 2018	November 2019
Updated the number of public health units to 35 due to the merger of Oxford, Elgin and St. Thomas public health units	November 2019
Added new data for 2017; re-extracted 2005–16 data and updated the tool	November 2018
Added LHIN data to interactive maps	March 2018
Added new data for 2016; re-extracted 2005–15 data and updated the tool	December 2017
Added interactive graphs and map for <i>Clostridium difficile</i> infection (CDI) outbreaks in hospitals	December 2017
Added interactive graphs to present 2016 specimen testing data for chlamydia, gonorrhea, influenza and legionellosis	December 2017
Added information on drug resistance for 2016 tuberculosis cases	December 2017

**Public Health Ontario**  
661 University Avenue, Suite 1701  
Toronto, Ontario  
M5G 1M1  
416.235.6556  
[communications@oahpp.ca](mailto:communications@oahpp.ca)  
[publichealthontario.ca](http://publichealthontario.ca)

