Organizational Risk Assessment for Clinical Office Practice

Public Health Ontario

Published: July 2024

Instructions

This template can be downloaded and used by clinical office practice to list risks for transmission of infections identified for each area of their clinic, analyze them, and to develop a plan to manage those risks.

For guidance on how to conduct an organizational risk assessment (ORA) and a sample, please

refer to the Infection Prevention and Control Orga	nizational Risk Assessment for Clinical Office
Practice available on PHO website.	
Area of the clinic:	
D / ODA	
Date ORA was conducted:	
Person(s) conducting ORA:	
1. First name:	Last name:
2. First name:	Last name.
2. First name:	Last name:
3. First name:	Last name:
Comments/additional information:	

Step 1. Identify Risks

Who could be exposed and how can they be exposed?

Step 2. Analyze the Risk

What could increase this risk of transmission and are there any control measures in place to prevent this transmission?	
Contributing factor(s):	
Control measure(s) in place:	
Step 3. Analyze the Risk	
What is the likelihood of this risk and the impact of transmission? Likelihood and impact of the identified risk (high, medium or low):	
Based on the above information select a level of risk:	
Risk Level and Actions: High (H): high likelihood and high severity - IMMEDIATE ACTION REQUIRED Medium (M): either likelihood or severity is high - Make efforts to further reduce the risk Low (L): low likelihood and low severity - Monitor and reduce the risk when feasible	

Step 4. Analyze the Risk

Can the risk be mitigated or reduced?
What additional control measures are needed?

Step 5. Implementation Plan

How will these additional control measures be implemented?
Who will be responsible?
Is there a way to know if these control measures work as expected?

Disclaimer:

This document may be freely used without permission for noncommercial purposes only and provided that appropriate credit is given to Public Health Ontario (PHO). No changes and / or modifications may be made to the content without express written permission from PHO.

For more information, visit publichealthontario.ca

